



Circumferential myocardial strain in cardiomyopathy with and without left bundle branch block

Citation

Han, Yuchi, Jonathan Chan, Idith Haber, Dana C Peters, Peter J Zimetbaum, Warren J Manning, and Susan B Yeon. 2010. Circumferential myocardial strain in cardiomyopathy with and without left bundle branch block. *Journal of Cardiovascular Magnetic Resonance* 12(1): 2.

Published Version

doi://10.1186/1532-429X-12-2

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:4728750>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Research

Open Access

Circumferential myocardial strain in cardiomyopathy with and without left bundle branch block

Yuchi Han^{*1}, Jonathan Chan^{1,2}, Idith Haber³, Dana C Peters¹, Peter J Zimetbaum¹, Warren J Manning^{1,4} and Susan B Yeon¹

Address: ¹Departments of Medicine (Cardiovascular Division), Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA, ²School of Medicine, University of Queensland, Brisbane, Australia, ³Department of Medicine (Cardiology), University of Chicago Medical School, Chicago, IL, USA and ⁴Department of Radiology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

Email: Yuchi Han^{*} - yhan@bidmc.harvard.edu; Jonathan Chan - j.chan2@uq.edu.au; Idith Haber - ihaber@medicine.bsd.uchicago.edu; Dana C Peters - dcpeters@bidmc.harvard.edu; Peter J Zimetbaum - pzimetba@bidmc.harvard.edu; Warren J Manning - wmanning@bidmc.harvard.edu; Susan B Yeon - syeon@bidmc.harvard.edu

^{*} Corresponding author

Published: 5 January 2010

Received: 4 August 2009

Journal of Cardiovascular Magnetic Resonance 2010, **12**:2 doi:10.1186/1532-429X-12-2

Accepted: 5 January 2010

This article is available from: <http://www.jcmr-online.com/content/12/1/2>

© 2010 Han et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Cardiac resynchronization therapy (CRT) has been shown to decrease mortality in 60-70% of advanced heart failure patients with left bundle branch block (LBBB) and QRS duration > 120 ms. There have been intense efforts to find reproducible non-invasive parameters to predict CRT response. We hypothesized that different left ventricular contraction patterns may exist in LBBB patients with depressed systolic function and applied tagged cardiovascular magnetic resonance (CMR) to assess circumferential strain in this population.

Methods: We determined myocardial circumferential strain at the basal, mid, and apical ventricular level in 35 subjects (10 with ischemic cardiomyopathy, 15 with non-ischemic cardiomyopathy, and 10 healthy controls). Patterns of circumferential strain were analyzed. Time to peak systolic circumferential strain in each of the 6 segments in all three ventricular slices and the standard deviation of time to peak strain in the basal and mid ventricular slices were determined.

Results: Dyskinesia of the anterior septum and the inferior septum in at least two ventricular levels was seen in 50% (5 out of 10) of LBBB patients while 30% had isolated dyskinesia of the antero-septum, and 20% had no dyskinesia in any segments, similar to all of the non-LBBB patients and healthy controls. Peak circumferential strain shortening was significantly reduced in all cardiomyopathy patients at the mid-ventricular level (LBBB $9 \pm 6\%$, non-LBBB $10 \pm 4\%$ vs. healthy $19 \pm 4\%$; both $p < 0.0001$ compared to healthy), but was similar among the LBBB and non-LBBB groups ($p = 0.20$). The LBBB group had significantly greater dyssynchrony compared to the non-LBBB group and healthy controls assessed by opposing wall delays and 12-segment standard deviation (LBBB 164 ± 30 ms vs. non-LBBB 70 ± 17 ms ($p < 0.0001$), non-LBBB vs. healthy 65 ± 17 ms ($p = 0.47$)).

Conclusions: Septal dyskinesia exists in some patients with LBBB. Myocardial circumferential strain analysis enables detailed characterization of contraction patterns, strengths, and timing in cardiomyopathy patients with and without LBBB.

Introduction

Cardiac resynchronization therapy (CRT) has been shown to improve symptoms, increase exercise capacity, decrease heart failure (HF) hospitalizations, and decrease mortality in patients with New York Heart Association (NYHA) Class III/IV HF with depressed systolic function, and a prolonged QRS in left bundle branch block (LBBB) morphology [1-3]. Recent data from patients with NYHA class I/II HF also demonstrated reduced HF hospitalization and reversal of left ventricular (LV) remodeling with CRT therapy [4]. However, 30-40% of patients who receive CRT therapy do not show significant clinical improvement [4-6]. As a result, there has been intense investigation to develop noninvasive parameters to predict CRT response [7-9]. While mechanical dyssynchrony assessed in the longitudinal axis of myocardial motion was shown to be predictive in single center trials [7-10], the multi-center PROSPECT trial failed to identify any echocardiographic dyssynchrony criteria to predict responders better than the clinical criteria [11].

Tagged cardiovascular magnetic resonance (CMR) is a noninvasive technique for measuring local deformation of the myocardium and quantitative assessment of mechanical dyssynchrony [12-14]. An advantage of tagged CMR circumferential strain (ϵ_{cc}) measurements is the narrow and consistent normal range across different centers [15,16]. In addition, ϵ_{cc} appears to be more sensitive to dyssynchrony than longitudinal strain in animal models [17]. ϵ_{cc} patterns in healthy patients have been studied in detail [16,18]. We sought to examine ϵ_{cc} patterns in patients with systolic dysfunction by applying tagged CMR.

Methods

Patient cohort

We studied twenty-five patients with systolic dysfunction referred for assessment of LV function and imaged between June 2006 and August 2009, including 10 patients with chronic ischemic cardiomyopathy (ICM) (age 64 ± 8 years, 90% male, LV ejection fraction (EF) $30 \pm 6\%$) and 15 patients with non-ischemic dilated cardiomyopathy (non-ICM) (age 59 ± 11 years, 73% male, LVEF $27 \pm 8\%$). All patients diagnosed with ICM had history of myocardial infarction and had coronary angiography demonstrating significant coronary artery disease involving at least two vessels. Eleven of 15 (73%) patients with non-ICM had coronary angiography demonstrating the absence of epicardial coronary artery stenoses. Four remaining patients were diagnosed as non-ICM with negative stress tests. Ten healthy adult subjects (age 38 ± 12 years, 50% male, EF $61 \pm 4\%$) served as controls. The institutional Committee on Clinical Investigation approved the study protocol. Written informed consent was

obtained from volunteers and was waived for existing clinical data sets.

ECG analysis

All subjects had a standard 12-lead ECG performed within a median of 15 days (with interquartile range of [5.5, 25.5] days) of the CMR with no intervening change in clinical status. The QRS morphology was determined by an experienced electrophysiologist (PJZ) according to AHA/ACCF/HRS guidelines [19]. Briefly, LBBB was determined if the QRS duration was ≥ 120 ms, with presence of a broad monophasic R wave in I, or V5 and V6, absence of Q waves in leads I, V5, and V6, and the displacement of the ST segment and T waves in a direction opposite to the major deflection of the QRS complex. The QRS duration was determined by automated computerized measurements and confirmed manually.

CMR

CMR studies were performed on a 1.5 T Philips Achieva MR scanner (Philips HealthCare, Best, NL), equipped with a 5-element cardiac coil. Breath-hold short-axis cine steady state free precession (SSFP) images covering the entire LV and long axis SSFP cine images covering the LV outflow tract were acquired as previously described [20].

Breath-hold ECG-gated tagged complementary spatial modulation of magnetization (CSPAMM) cine images at the basal, mid, and apical ventricular levels were obtained [21,22]. The mid ventricular level was prescribed at the mid-papillary muscle level. The center of the basal slice and the center of the apical slice were acquired 20 mm proximal and distal to the mid slice center, respectively. Scan parameters include spiral readout with 8 interleaves, 9 ms acquisition window, repetition time (TR)/echo time (TE)/flip angle (α) = 25 ms/3.6 ms/25°, field of view (FOV) = 320 mm \times 320 mm, 10 mm slice thickness with 5 mm tag spacing, temporal resolution 25-35 ms, spatial resolution 2.5 \times 2.5 \times 10 mm.

Free-breathing, ECG-triggered phase contrast velocity sequences for aortic flow oriented in the axial plane at the level of the bifurcation of the pulmonary artery were acquired as previously described [23]. Sequence parameters were: TR/TE/ α = 15 ms/6.5 ms/30°, FOV = 300 mm \times 210 mm, matrix = 128 \times 128, slice thickness 6 mm. Respiratory motion compensation was accomplished with the use of three signal averages.

2D breath-hold ECG-triggered late gadolinium enhancement (LGE) images were acquired in the same orientation as SSFP short axis images and long axis 2-chamber and 4-chamber orientations at 10-20 minutes post injection of 0.2 mmol/kg gadolinium-diethylenetriamine pentaacetic acid (Magnevist, Schering, Germany). Imaging parameters

were: 2D spoiled gradient echo inversion recovery, TR/TE/ $\alpha = 4.3$ ms/1.5 ms/20°, FOV = 320 mm × 320 mm, matrix = 160 × 160, 8 mm slices with 2 mm gaps, partial echo, fat saturation, 1 RR between inversions, and two signal averages.

Volumetric Analysis

Cardiac volumes were calculated in the standard fashion as previously described with papillary muscle included in the LV cavity volume [20]. Mitral regurgitation volume = LV stroke volume - aortic forward flow volume.

Timing of systole

Systolic ejection begins when the aortic valve opens, as seen from the cine long axis LV outflow tract images and

confirmed with phase contrast aortic flow curves. End of systole is defined as the time of aortic valve closure.

CSPAMM image analysis

A customized software program (Cardiotool), written in MATLAB (MathWorks, Natick, MA), was used for semi-automated analysis of circumferential strain [24]. Endocardial and epicardial borders were drawn manually on the tagged images, and the right ventricular insertion sites were marked to indicate the outer borders of the antero-septum and inferoseptum (Figure 1). The remaining myocardial slice was divided into anterior, anterolateral, inferolateral, and inferior segments according to the AHA 17-segment model. Circumferential strain from the mid-myocardial layer of each of the six segments of all three ventricular slices was analyzed.




Figure 1

CSPAMM images of mid ventricular short axis slice and analysis model. A. Tags at the beginning of systole after ECG triggering. B. Same slice at end systole. C. Same slice at diastole with good persistence of tags. D and E show an example of CSPAMM images with the analysis model. D. In early systole, the epicardial border was manually drawn in green, endocardial border in red, superior right ventricular insertion site marked with a yellow triangle, and inferior insertion site marked with a blue triangle. The red donut is the analysis model with the myocardium partitioned in epi-myocardial, mid-myocardial, and endo-myocardial thirds shown in early systole in D and end-systole in E. Magenta dot is the center of the epicardial contour.

LGE image analysis

The presence of abnormal LGE in the myocardium was determined by the presence of high signal intensity (defined as 6 standard deviations higher than remote myocardium) in the short axis images and confirmed in the long axis images. The degree of LGE was determined by <50%, 50-75%, and >75% of LGE transmural.

Dyssynchrony assessment

Time to peak myocardial systolic strain in the mid-myocardial layer was identified for each segment by identifying the cardiac phase of peak systolic strain and multiplying by the temporal resolution in milliseconds. In myocardial segments with presystolic negative ϵ_{cc} and systolic positive ϵ_{cc} , the peak time was taken at the presystolic negative ϵ_{cc} . For other segments, the peak time was taken at the maximal circumferential shortening, including the post-systolic shortening period. The absolute time difference between opposing walls (inferolateral wall to anteroseptum (IL-AS), anterior wall to inferior wall (A-I), and anterolateral wall to inferoseptum (AL-IS)) were obtained. The standard deviation of time to peak systolic strain in the basal and mid segments (T12SD) was determined.

Statistics

Data were analyzed using the two-tailed Student's t-test to compare continuous variables and the Wilcoxon rank sum test to compare categorical variables. A two tailed p-value of < 0.05 was considered significant. In multiple group comparisons, Bonferroni correction was applied. All statistical analyses were performed with STATA Version 10 (STATcorp, TX, USA).

Results**Patient characteristics**

Patient clinical characteristics are presented in Table 1. Twelve (48%) patients with ICM ($n = 4$) and non-ICM ($n = 8$) had LBBB with a mean QRS duration of 161 ± 10 ms. The thirteen non-LBBB patients had either normal QRS duration (< 100 ms) ($n = 6$) of 81 ± 5 ms or interventricular conduction delay (IVCD) ($n = 7$) with a mean QRS duration of 118 ± 14 ms. The healthy control subjects had normal QRS morphology and a mean QRS duration of 88 ± 12 ms.

The LBBB and non-LBBB patient groups had similar age, gender, prevalence of hypertension, diabetes, hyperlipidemia, tobacco use, cardiac medication use, LVEF, LV volumes, and mitral regurgitation volume, but differed in their NYHA classification (Table 1).

Myocardial scar

Among the four ICM patients with LBBB, three had inferior infarcts (two with LGE > 75%, one had LGE <50%)

and one patient had both inferior and anterior infarcts (LGE = 50% of the septum, anterior and inferior walls). Among the six ICM patients with non-LBBB, four had anterior infarcts (3 subendocardial LGE, and one with > 75% of anteroseptum and anterior wall LGE) and two had inferior infarcts (one subendocardial LGE, one >75% LGE). No focal LGE was identified in patients with non-ICM.

Circumferential strain patterns in LBBB

Three distinctive contractile patterns in LBBB patients were observed in the septum.

Type I: An initial negative ϵ_{cc} in the septal segment before systolic ejection (as determined above) was present as presystolic contraction, followed by a positive ϵ_{cc} reflecting stretching and dyskinesis of the septum. We further divided this type into two subtypes:

Type Ia. When Type I pattern is only present in the anteroseptum (Figure 2A).

Type Ib. When Type I septal pattern is present in both anteroseptum and inferoseptum (Figure 2C).

Type II. Decreased amplitude but no stretching in either of the septal segments (Figure 2E).

One LBBB patient had no basal or apical ventricular CSPAMM imaging. One patient was excluded from pattern analysis due to poor basal slice image quality. For the remaining 10 patients, 6 patients had fully concordant contractile pattern in the septum in all three slices. Two had basal and mid slice concordance and two had mid and apical slice concordance. Type II pattern was more prevalent in apical slices (50%) compared to 20% of mid slices and 10% of basal slices. Overall, the mid ventricular slice was the most representative of septal contraction pattern with 100% concordance with either the basal or apical slice or both. Among LBBB patients, 30% of had Type Ia, 50% of had Type Ib while 20% had Type II in the mid ventricular slice. These two Type II patients had QRS durations of 156 ms and 180 ms respectively. The different strain patterns cannot be predicted from ECG pattern or QRS duration in these LBBB patients (Figure 2B, D and 2F).

In addition, Type I pattern was present in the anterior segment in the mid slice in two of the three Type Ia patients and two of the five Type Ib patients. All remaining segments in all three slices had negative ϵ_{cc} during systole.

Table 1: LBBB and non-LBBB patient characteristics:

| | LBBB (n = 12) | Non-LBBB (n = 13) | p value |
|---------------------------|----------------------|--------------------------|----------------|
| Age (years) | 60 ± 10 | 62 ± 11 | 0.729 |
| Sex (% male) | 83% | 77% | 0.702 |
| % ICM | 33% | 46% | 0.532 |
| LVEF (%) | 27 ± 7 | 30 ± 7 | 0.327 |
| LVEDV (ml) | 307 ± 60 | 264 ± 60 | 0.086 |
| LVESV (ml) | 228 ± 56 | 189 ± 62 | 0.115 |
| Mitral Regurgitation (ml) | 7 ± 12 | 8 ± 8 | 0.817 |
| QRS Duration (ms) | 161 ± 10 | 106 ± 16* | < 0.0001 |
| NYHA class (median) | 2.5 | 1* | 0.003 |
| Hypertension | 67% | 85% | 0.320 |
| Diabetes | 25% | 31% | 0.760 |
| Hyperlipidemia | 33% | 69% | 0.078 |
| Tobacco Use | 42% | 46% | 0.830 |
| ACEI/ARB | 100% | 92% | 0.337 |
| Beta-blocker | 100% | 85% | 0.165 |
| Diuretic | 67% | 31% | 0.078 |
| Aspirin | 67% | 77% | 0.589 |
| Digoxin | 42% | 8% | 0.059 |
| Spironolactone | 25% | 15% | 0.571 |

± standard deviation. ICM = ischemic cardiomyopathy. LVEF = left ventricular ejection fraction. LVEDV = left ventricular end-diastolic volume. LVESV = left ventricular end-systolic volume. NYHA = New York Heart Association Classification. ACEI/ARB = angiotensin converting enzyme inhibitor/angiotensin receptor blocker. *p < 0.05.


Circumferential strain pattern in patients without LBBB and healthy subjects

All patients without LBBB and all healthy subjects showed ϵ_{cc} shortening in all segments throughout systolic contraction (Figure 3) in all slices at the basal, mid, and apical ventricular levels. The septal contractile pattern was similar in these subjects.


Circumferential shortening patterns in ischemic vs. non-ischemic groups

We found no difference in time to peak ϵ_{cc} in ischemic and non-ischemic patients stratified by LBBB and non-LBBB.


In three out of four patients with evidence for > 75% scar in the infarcted walls, there was no circumferential contraction ($\epsilon_{cc} \approx 0$). These three segments were excluded from timing analysis due to the absence of peak circumferential shortening. All remaining infarct segments were included in the analysis. One patient with ICM and non-LBBB had > 75% scar in the anteroseptum, but the ϵ_{cc} in that segment was not near 0. In Figure 4, we show side by side the ϵ_{cc} in the anteroseptum and inferoseptum of ICM and non-ICM patients with Type Ia, Type Ib, and Type II LBBB patterns.

**Figure 2**

Three different septal circumferential strain patterns in patients with LBBB. AVO marks the opening of aortic valve and AVC marks the closure of aortic valve. A. Type Ia. Dyskinesia of the anteroseptum. Inferoseptum ϵ_{cc} shortening is reduced. B. Corresponding ECG, QRS = 154 ms. C. Type Ib. Dyskinesia of the entire septum. D. Corresponding ECG, QRS = 150 ms. E. Type II. No dyskinesia but reduced septal ϵ_{cc} . F. Corresponding ECG, QRS = 156 ms. Arrows point to presystolic contraction. AS = anteroseptum, IS = inferoseptum, and IL = inferolateral wall. AVO = aortic valve opening. AVC = aortic valve closure.

**Figure 3**

Septal circumferential strain pattern in patients with non-LBBB (A, B) and healthy (C) subjects. All subjects had normal contraction patterns with maximum negative circumferential strain reached in the septum earlier than in the inferolateral wall. A. Patient with interventricular conduction delay and QRS duration of 146 ms. B. Patient with a normal QRS duration of 98 ms. C. Healthy subject with a normal QRS duration of 98 ms. AS = anteroseptum, IS = inferoseptum, IL = inferolateral wall. AVO = aortic valve opening. AVC = aortic valve closure.

**Figure 4**

All three types of septal contraction pattern can be observed in both ischemic (ICM) and non-ischemic (non-ICM) cardiomyopathy patients. In LBBB Type Ia, only the anteroseptum is dyskinetic. In LBBB Type Ib, both anteroseptum and inferoseptum are dyskinetic. In LBBB Type II, neither anteroseptum nor inferoseptum is dyskinetic. AS = anteroseptum, IS = inferoseptum.

Circumferential shortening

A total of 24 of 618 segments (4%) from all subjects and all slices were excluded from analysis due to poor image quality. Cardiomyopathy patients showed reduced overall circumferential shortening (%S) in the mid slice (LBBB $9 \pm 6\%$, non-LBBB $10 \pm 4\%$ vs. healthy $19 \pm 4\%$) (both $p < 0.0001$ compared to healthy) and apical slice (LBBB $8 \pm 6\%$, non-LBBB $10 \pm 5\%$ vs. healthy $19 \pm 5\%$) (both $p < 0.0001$ compared to healthy) (Figure 5). Within the mid slice, in the non-LBBB group, %S was reduced in all segments compared to healthy controls (all $p < 0.0001$); while in the LBBB group, %S was reduced in all segments except the anterolateral wall as compared to controls. The LBBB patients had a significant reduction in %S of ϵ_{cc} in the anteroseptum ($3 \pm 2\%$ vs. $9 \pm 4\%$) ($p = 0.001$) and anterior segment ($6 \pm 4\%$ vs. $10 \pm 2\%$) ($p = 0.009$) compared to non-LBBB patients, but not overall ($p = 0.20$) (Table 2).

Dyssynchrony

In patients with LBBB compared to non-LBBB or healthy subjects, there was significantly greater dyssynchrony of contraction as indicated by greater delays in the opposing

walls IL-AS and AL-IS at all three ventricular levels, and T12SD ($p \leq 0.002$) (Table 3). When comparing non-LBBB to healthy subjects, only basal IL-AS delay and mid AL-IS delay were significantly lengthened ($p \leq 0.002$). The A-I time to peak strain had no significant delays in all three groups.

Interobserver and intraobserver variability

In six randomly selected subjects with 35 evaluable segments in the mid-ventricular slice, intraobserver variability and interobserver variability for peak systolic ϵ_{cc} timing were $3 \pm 7\%$ and $7 \pm 11\%$, respectively. The intraobserver variability and interobserver variability for systolic ϵ_{cc} %S were $9 \pm 8\%$, $9 \pm 9\%$, respectively. Both observers on repeated analysis identified the same contractile patterns.

Discussion

In our study of patients with LBBB pattern and depressed LV function, we found three types of septal myocardial circumferential strain patterns. Some patients with LBBB had severe mechanical dyssynchrony manifested as a specific contractile pattern with initial presystolic septal contraction during isovolumic contraction period followed by


Table 2: Comparison of circumferential myocardial strain percent shortening of all three ventricular slices in patients with and without LBBB, and healthy subjects.

| | | LBBB (%S) | Non-LBBB (%S) | Healthy (%S) | p(LBBB vs. non LBBB)* | p(LBBB vs. healthy)* | p(non-LBBB vs. healthy)* |
|--------|----|-----------|---------------|--------------|-----------------------|----------------------|--------------------------|
| Basal | AS | 3 ± 2 | 6 ± 3 | 13 ± 4 | 0.005* | < 0.0001* | 0.0009* |
| | A | 9 ± 4 | 8 ± 4 | 11 ± 5 | 0.624 | 0.285 | 0.115 |
| | AL | 17 ± 3 | 15 ± 2 | 14 ± 3 | 0.121 | 0.066 | 0.502 |
| | IL | 16 ± 6 | 11 ± 4 | 11 ± 3 | 0.046 | 0.027 | 0.683 |
| | I | 9 ± 6 | 7 ± 3 | 8 ± 2 | 0.371 | 0.806 | 0.296 |
| | IS | 4 ± 2 | 6 ± 4 | 8 ± 3 | 0.177 | 0.003* | 0.191 |
| Mid | AS | 3 ± 2 | 9 ± 4 | 18 ± 2 | 0.001* | < 0.0001* | < 0.0001* |
| | A | 6 ± 4 | 10 ± 2 | 20 ± 3 | 0.009* | < 0.0001* | < 0.0001* |
| | AL | 17 ± 4 | 14 ± 2 | 20 ± 3 | 0.071 | 0.047 | = 0.0004* |
| | IL | 16 ± 4 | 12 ± 4 | 24 ± 3 | 0.026 | < 0.0001* | < 0.0001* |
| | I | 7 ± 3 | 7 ± 5 | 16 ± 4 | 0.823 | < 0.0001* | 0.0003* |
| | IS | 4 ± 3 | 7 ± 4 | 17 ± 3 | 0.036 | < 0.0001* | < 0.0001* |
| Apical | AS | 5 ± 4 | 9 ± 4 | 18 ± 3 | 0.012* | < 0.0001* | < 0.0001* |
| | A | 6 ± 5 | 12 ± 5 | 18 ± 4 | 0.007* | < 0.0001* | 0.004* |
| | AL | 14 ± 4 | 14 ± 3 | 23 ± 5 | 0.891 | 0.0003* | 0.0003* |
| | IL | 16 ± 4 | 13 ± 4 | 22 ± 7 | 0.096 | 0.022 | 0.002* |
| | I | 6 ± 3 | 7 ± 5 | 17 ± 3 | 0.565 | < 0.0001* | < 0.0001* |
| | IS | 5 ± 4 | 8 ± 5 | 16 ± 2 | 0.095 | < 0.0001* | 0.0001* |

± standard deviation. *with Bonferroni correction for multiple comparisons, $p < 0.017$ was considered significant. AS = antero-septum. A = anterior wall. AL = anterolateral wall. IL = inferolateral wall. I = inferior wall. IS = inferoseptum.

dyskinesia (positive ϵ_{cc}) of the interventricular septum during the entire systole. This pattern was present in the antero-septum (Type Ia) in 30% of patients, and in the entire septum in 50% of patients (Type Ib). The remaining 20% of LBBB patients had a normal contractile pattern, similar to non-LBBB cardiomyopathy patients and healthy controls, although the magnitude of contraction is significantly reduced in both groups of cardiomyopathy patients compared to healthy controls. All non-septal segments except four anterior segments in LBBB patients demonstrated ϵ_{cc} shortening. Our finding of ϵ_{cc} of 16-24% in normal healthy controls is consistent with literature findings in the mid-myocardial layer [15].

Detailed studies of healthy human subjects have demonstrated that the normal mechanical activation pattern starts at the septum and extends to the inferolateral wall [18]. We found presystolic mid-myocardial circumferential shortening (negative ϵ_{cc}) during isovolumic contraction phase in part or the entire septum in patients with LBBB, followed by circumferential lengthening (positive ϵ_{cc}) during systole. This altered pattern with early presystolic contraction and systolic dyskinesia contribute to increased severity of dyssynchrony. In our cohort of ICM and non-ICM patients, the presence or absence of LBBB had a dominating impact on the contractile pattern regardless of the etiology of the cardiomyopathy. In ICM

**Figure 5**

Circumferential shortening magnitude in three ventricular slices. Compared to healthy controls, both the LBBB and non-LBBB patients have overall significantly reduced ϵ_{cc} %S in mid and apical ventricular slices. In the basal slice, non-LBBB patients had significantly reduced ϵ_{cc} %S compared to healthy controls but not the LBBB patients. %S = % shortening.

patients, the only difference was the presence of akinesis in regions of transmural infarction in 3 segments. Our data for circumferential strain demonstrate that there are significant timing differences in opposing walls (IL-AS, AL-IS), and T12SD in patients with low EF and LBBB as compared to the patients with low EF but normal QRS or IVCD. The prognostic clinical significance of these dyssynchrony patterns to CRT remains to be determined.

In a 2D longitudinal strain by speckle-tracking study by Carasso et al, dyssynchrony timing was not the only predictor for CRT success [25]. In their cohort, the majority of non-responders to CRT had ≥ 1 segment of "holosystolic stretching" (dyskinesia) while all except one responder had no such segments [25]. In a non-responder, the holosystolic dyskinetic segment remained dyskinetic after CRT; in a responder, the early systolic dyskinetic segment had normal contractile pattern after CRT therapy [25]. Our study also demonstrates the existence of different pattern of contraction within LBBB, which may be as important as timing in the assessment of dyssynchrony in patients with systolic dysfunction. Further investigation is needed to determine whether the septal contractile patterns impact response to CRT.

This study has several limitations. We did not study patients with normal EF and LBBB, low EF and RBBB, or a low EF and narrow LBBB (120-149 ms). In a 2D speckle

tracking study by Miyazaki et al, patients with LBBB and normal EF had increased opposing wall delays and increased standard deviation of time to peak longitudinal strain as compared to normal controls, but the increase was less than patients with reduced EF and normal QRS [26]. The patients with reduced EF < 35% and LBBB had the longest delays [26]. Our findings on circumferential strain in low EF and LBBB vs. non-LBBB patients are consistent with their results. Patients with different conduction abnormalities would be important to study in order to further define and understand the different mechanical contractile patterns associated with conduction abnormalities.

Conclusions

We have demonstrated LBBB in some patients with systolic dysfunction is associated with dyskinesia of the antero-septum or the entire septum, resulting in severe mechanical dyssynchrony. The strength of myocardial contraction is significantly reduced in cardiomyopathic patients regardless of conduction patterns. The recognition of the presence of different mechanical contraction patterns within the same conduction abnormality may be important for the selection of patients for CRT.

List of abbreviations

CMR: cardiovascular magnetic resonance; CRT: Cardiac resynchronization therapy; HF: heart failure; LV: left ven-

Table 3: Comparison of dyssynchrony measurements in three ventricular slices in patients with and without LBBB, and healthy subjects.

| | | LBBB | Non-LBBB | Healthy | p(LBBB vs. non LBBB)* | p(LBBB vs. healthy)* | p(non-LBBB vs. healthy)* |
|------------|-----------------------|---------------------|----------------------|--------------------|------------------------------|-----------------------------|---------------------------------|
| Basal | IL-AS (ms) (range) | 391 ± 37 (325-425) | 136 ± 79 (0-325) | 45 ± 23 (0-75) | < 0.0001* | < 0.0001* | 0.001* |
| | AL-IS (ms) (range) | 284 ± 119 (150-425) | 106 ± 54 (0-200) | 68 ± 64 (0-175) | 0.001* | 0.0002* | 0.171 |
| | A-I (ms) (range) | 122 ± 145 (0-400) | 83 ± 85 (0-225) | 68 ± 32 (0-100) | 0.478 | 0.272 | 0.594 |
| Mid | IL-AS (ms) (range) | 353 ± 79 (245-425) | 82 ± 44 (25-140) | 69 ± 39 (0-125) | < 0.0001* | < 0.0001* | 0.451 |
| | AL-IS (ms) (range) | 283 ± 108 (125-400) | 114 ± 37 (25-175) | 50 ± 46 (0-125) | 0.0002* | < 0.0001* | 0.002* |
| | A-I (ms) (range) | 168 ± 164 (0-475) | 69 ± 46 (0-175) | 64 ± 55 (0-150) | 0.097 | 0.084 | 0.791 |
| Apical | IL-AS (ms) (range) | 278 ± 118 (140-425) | 61 ± 47 (0-325) | 38 ± 41 (0-75) | < 0.0001* | < 0.0001* | 0.260 |
| | AL-IS (ms) (range) | 224 ± 118 (150-425) | 70 ± 63 (0-200) | 54 ± 24 (0-175) | 0.002* | 0.001* | 0.437 |
| | A-I (ms) (range) | 96 ± 110 (0-300) | 48 ± 43 (0-150) | 34 ± 20 (0-75) | 0.200 | 0.093 | 0.601 |
| T12SD (ms) | | 164 ± 30 (131-207) | 70 ± 17 (51-103) | 65 ± 17 (44-97) | < 0.0001* | < 0.0001* | 0.467 |

Data presented as absolute time difference between the two walls ± standard deviation. *with Bonferroni correction for multiple comparisons, $p < 0.017$ was considered significant. IL-AS = inferolateral wall to anteroseptal wall. AL-IS = anterolateral wall to inferoseptal wall. A-I = Anterior wall to inferior wall.

tricle; LBBB: left bundle branch block; NYHA: New York Heart Association; ICM: ischemic cardiomyopathy; SSFP: steady state free precession; TE: echo time; TR: repetition time; FOV: field of view; LGE: late gadolinium enhancement; CSPAMM: complementary spatial modulation of magnetization; IL-AS: inferolateral to anteroseptal; AL-IS: anterolateral to inferoseptal; A-I: anterior to inferior; IVCD: interventricular conduction delay; T12SD: Standard deviation of time to peak systole circumferential strain in the 12 segments of basal and mid ventricular slices; %S: percent shortening.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

YH designed the study, acquired data, analyzed and interpreted the data, and drafted the manuscript. JC analyzed and interpreted the data and critically revised the manuscript. IH and DCP participated in the analysis and interpretation of the data, and critical review of the manuscript. PJZ participated in the conception and design of study and ECG analysis. WJM participated in interpretation

and critical review of the manuscript. SBY conceived the study, participated in the design, interpretation, and critical review of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We like to acknowledge Kraig V. Kissinger, MRT and Beth Goddu, MRT for their assistance in data collection.

Grant Support:

American College of Cardiology Foundation/GE Career Development Award in Cardiovascular Imaging (to YH);

Beth Israel Deaconess Medical Center and the Clinical Investigator Training Program: Beth Israel Deaconess Medical Center - Harvard/MIT Health Sciences and Technology, in collaboration with Pfizer Inc. and Merck & Co. (to YH);

Royal Australasian College of Physicians (to JC);

American Heart Association Scientist Development Award (to DCP);

NIH NIBIB Career Research Award K01 (to DCP).

All funding sources provide protected time for the investigators and are not involved in the study design, data collection, analysis, and interpretation, or the writing and submission of the manuscript.

References

- Cazeau S, Leclercq C, Lavergne T, Walker S, Varma C, Linde C, Garrigue S, Kappenberger L, Haywood GA, Santini M, Bailleul C, Daubert JC, Multisite Stimulation in Cardiomyopathies (MUSTIC) Study Investigators: **Effects of multisite biventricular pacing in patients with heart failure and intraventricular conduction delay.** *N Engl J Med* 2001, **344**:873-880.
- Bradley DJ, Bradley EA, Baughman KL, Berger RD, Calkins H, Goodman SN, Kass DA, Powe NR: **Cardiac resynchronization and death from progressive heart failure: a meta-analysis of randomized controlled trials.** *Jama* 2003, **289**:730-740.
- Cleland JG, Daubert JC, Erdmann E, Freemantle N, Gras D, Kappenberger L, Tavazzi L: **The effect of cardiac resynchronization on morbidity and mortality in heart failure.** *N Engl J Med* 2005, **352**:1539-1549.
- Linde C, Abraham WT, Gold MR, St John Sutton M, Ghio S, Daubert C: **Randomized trial of cardiac resynchronization in mildly symptomatic heart failure patients and in asymptomatic patients with left ventricular dysfunction and previous heart failure symptoms.** *J Am Coll Cardiol* 2008, **52**:1834-1843.
- Abraham WT, Fisher WG, Smith AL, Delurgio DB, Leon AR, Loh E, Kocovic DZ, Packer M, Clavell AL, Hayes DL, Ellestad M, Trupp RJ, Underwood J, Pickering F, Truex C, McAtee P, Messenger J, MIRACLE Study Group: **Multicenter InSync Randomized Clinical Evaluation: Cardiac resynchronization in chronic heart failure.** *N Engl J Med* 2002, **346**:1845-1853.
- Bristow MR, Saxon LA, Boehmer J, Krueger S, Kass DA, De Marco T, Carson P, DiCarlo L, DeMets D, White BG, DeVries DW, Feldman AM, Comparison of Medical Therapy, Pacing, and Defibrillation in Heart Failure (COMPANION) Investigators: **Cardiac-resynchronization therapy with or without an implantable defibrillator in advanced chronic heart failure.** *N Engl J Med* 2004, **350**:2140-2150.
- Yu CM, Zhang Q, Fung JW, Chan HC, Chan YS, Yip GW, Kong SL, Lin H, Zhang Y, Sanderson JE: **A novel tool to assess systolic asynchrony and identify responders of cardiac resynchronization therapy by tissue synchronization imaging.** *J Am Coll Cardiol* 2005, **45**:677-684.
- Penicka M, Bartunek J, De Bruyne B, Vanderheyden M, Goethals M, De Zutter M, Brugada P, Geelen P: **Improvement of left ventricular function after cardiac resynchronization therapy is predicted by tissue Doppler imaging echocardiography.** *Circulation* 2004, **109**:978-983.
- Bax JJ, Bleeker GB, Marwick TH, Molhoek SG, Boersma E, Steendijk P, Wall EE van der, Schalij MJ: **Left ventricular dyssynchrony predicts response and prognosis after cardiac resynchronization therapy.** *J Am Coll Cardiol* 2004, **44**:1834-1840.
- Yu CM, Chau E, Sanderson JE, Fan K, Tang MO, Fung WH, Lin H, Kong SL, Lam YM, Hill MR, Lau CP: **Tissue Doppler echocardiographic evidence of reverse remodeling and improved synchronicity by simultaneously delaying regional contraction after biventricular pacing therapy in heart failure.** *Circulation* 2002, **105**:438-445.
- Chung ES, Leon AR, Tavazzi L, Sun JP, Nihoyannopoulos P, Merlino J, Abraham WT, Ghio S, Leclercq C, Bax JJ, Yu CM, Gorcsan J, St John Sutton M, De Sutter J, Murillo J: **Results of the Predictors of Response to CRT (PROSPECT) trial.** *Circulation* 2008, **117**:2608-2616.
- Lardo AC, Abraham TP, Kass DA: **Magnetic resonance imaging assessment of ventricular dyssynchrony: current and emerging concepts.** *J Am Coll Cardiol* 2005, **46**:2223-2228.
- Gotte MJ, Germans T, Russel IK, Zwanenburg JJ, Marcus JT, van Rossum AC, van Veldhuisen DJ: **Myocardial strain and torsion quantified by cardiovascular magnetic resonance tissue tagging: studies in normal and impaired left ventricular function.** *J Am Coll Cardiol* 2006, **48**:2002-2011.
- Bilchick KC, Dimaano V, Wu KC, Helm RH, Weiss RG, Lima JA, Berger RD, Tomaselli GF, Bluemke DA, Halperin HR, Abraham T, Kass DA, Lardo AC: **Cardiac magnetic resonance assessment of dyssynchrony and myocardial scar predicts function class improvement following cardiac resynchronization therapy.** *JACC Cardiovasc Imaging* 2008, **1**:561-568.
- Moore CC, McVeigh ER, Zerhouni EA: **Quantitative tagged magnetic resonance imaging of the normal human left ventricle.** *Top Magn Reson Imaging* 2000, **11**:359-371.
- Moore CC, Lugo-Olivieri CH, McVeigh ER, Zerhouni EA: **Three-dimensional systolic strain patterns in the normal human left ventricle: characterization with tagged MR imaging.** *Radiology* 2000, **214**:453-466.
- Helm RH, Leclercq C, Faris OP, Ozturk C, McVeigh E, Lardo AC, Kass DA: **Cardiac dyssynchrony analysis using circumferential versus longitudinal strain: implications for assessing cardiac resynchronization.** *Circulation* 2005, **111**:2760-2767.
- Zwanenburg JJ, Gotte MJ, Kuijjer JP, Heethaar RM, van Rossum AC, Marcus JT: **Timing of cardiac contraction in humans mapped by high-temporal-resolution MRI tagging: early onset and late peak of shortening in lateral wall.** *Am J Physiol Heart Circ Physiol* 2004, **286**:H1872-1880.
- Surawicz B, Childers R, Deal BJ, Gettes LS, Bailey JJ, Gorgels A, Hancock EW, Josephson M, Kligfield P, Kors JA, Macfarlane P, Mason JW, Mirvis DM, Okin P, Pahlm O, Rautaharju PM, van Herpen G, Wagner GS, Wellens H, American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology; American College of Cardiology Foundation; Heart Rhythm Society: **AHA/ACCF/HRS recommendations for the standardization and interpretation of the electrocardiogram: part III: intraventricular conduction disturbances: a scientific statement from the American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology; the American College of Cardiology Foundation; and the Heart Rhythm Society: endorsed by the International Society for Computerized Electrocardiology.** *Circulation* 2009, **119**:235-240.
- Salton CJ, Chuang ML, O'Donnell CJ, Kupka MJ, Larson MG, Kissinger KV, Edelman RR, Levy D, Manning WJ: **Gender differences and normal left ventricular anatomy in an adult population free of hypertension. A cardiovascular magnetic resonance study of the Framingham Heart Study Offspring cohort.** *J Am Coll Cardiol* 2002, **39**:1055-1060.
- Haber I, Metaxas DN, Axel L: **Three-dimensional motion reconstruction and analysis of the right ventricle using tagged MRI.** *Med Image Anal* 2000, **4**:335-355.
- Ryf S, Kissinger KV, Spiegel MA, Bornert P, Manning WJ, Boesiger P, Stuber M: **Spiral MR myocardial tagging.** *Magn Reson Med* 2004, **51**:237-242.
- Gelfand EV, Hughes S, Hauser TH, Yeon SB, Goepfert L, Kissinger KV, Rofsky NM, Manning WJ: **Severity of mitral and aortic regurgitation as assessed by cardiovascular magnetic resonance: optimizing correlation with Doppler echocardiography.** *J Cardiovasc Magn Reson* 2006, **8**:503-507.
- Haber I, Kikinis R, Westin C-F: **Phase-Driven Finite Element Model for Spatio-temporal Tracking in Cardiac Tagged MRI.** *Medical Image Computing and Computer-Assisted Intervention* 2001 2001:1332-1335.
- Carasso S, Rakowski H, Witte KK, Smith P, Carasso D, Garceau P, Sasson Z, Parker JD: **Left ventricular strain patterns in dilated cardiomyopathy predict response to cardiac resynchronization therapy: timing is not everything.** *J Am Soc Echocardiogr* 2009, **22**:242-250.
- Miyazaki C, Powell BD, Bruce CJ, Espinosa RE, Redfield MM, Miller FA, Hayes DL, Cha YM, Oh JK: **Comparison of echocardiographic dyssynchrony assessment by tissue velocity and strain imaging in subjects with or without systolic dysfunction and with or without left bundle-branch block.** *Circulation* 2008, **117**:2617-2625.